

The Hawking-Unruh phenomenon on graphene

Alfredo Iorio¹ & Gaetano Lambiase^{2,3}

¹*Faculty of Mathematics and Physics, Charles University in Prague, V Holešovičkách 2, 18000 Prague 8 - Czech Republic.*

²*Department of Physics, University of Salerno, Fisciano (SA) - Italy.*

³*INFN, Sezione di Napoli - Italy.*

A central issue in the ongoing studies of graphene is how the morphology of the sample modifies its electronic properties. For curved shapes its well-known special relativistic-like properties naturally point to quantum field theory in curved spaces that is not a general relativistic-like description as yet because the time components of the metric are left over. Here we find that, if graphene is shaped as Beltrami pseudosphere the problem is solved. As a result, through the appropriate identifications of the graphene counterpart of the customary relativistic procedures, we are able to predict that the curved graphene sample has a finite temperature electronic local density of states that is a realization of the Hawking-Unruh effect. We then propose an experiment with a Scanning Tunneling Microscope that should easily detect such effect and measure the Hawking temperature.

1 Graphene and quantum field theory in curved spacetimes

Graphene is an allotrope of carbon that was first theoretically posited^{1,2} and then found to have an abundance of “unorthodox” properties^{3,4}, the understanding of which is appealing to theorists

and experimentalists, see, e.g., ⁵. Following the spirit of the condensed matter analogues of gravitational effects^{6,7} we shall use graphene as a physical realization of quantum field theory (QFT) in curved spacetimes⁸⁻¹¹. Of these unorthodox properties one that is crucial is the dimensionality (we shall evoke the exotic scenarios of gravity in 2+1 dimensions^{12,13}) and another one is the well known massless Dirac field description of graphene's conductivity electrons¹⁴ (we shall use the fact that the theory enjoys a scale symmetry proposed by Weyl in an early attempt to introduce the gauge principle^{15,16}).

In this work we assume that the conductivity electrons of graphene experience a general relativistic-like spacetime and we identify a specific shape, the Beltrami pseudosphere, for which it is easier to probe this assumption. In turn, the predictions of QFT in curved spacetimes point to a thermal spectrum in the form of a finite temperature electronic local density of states (LDOS). The temperature is of the Hawking-Unruh type ^{17,18}. It is of geometric origin and depends upon the curvature and upon the the meridian coordinate on the surface. The final outcome of this study is an exact formula for the LDOS that we show to be possible to measure. The whole construction, including the assumption on graphene's electrons experiencing a curved spacetime, can be submitted to the judgement of experiments. Among the unusual features of the formula is a Bose-Einstein thermal spectrum for this Dirac system that is due to the odd dimensions^{10,19,20}.

2 Shaping graphene as a Beltrami pseudosphere and the issue of time

For graphene, to include time in a relativistic fashion we need to take two steps. First, we have to assume that for a curved graphene sample the conductivity electrons experience a spacetime metric, the choice dictated by everyday practice being

$$g_{\mu\nu}^{\text{graphene}}(q) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & & \\ & g_{\alpha\beta} & \\ 0 & & \end{pmatrix}, \quad (1)$$

i.e., the time-time component is just the flat one. Here $\mu, \nu = 0, 1, 2$ and $\alpha, \beta = 1, 2$, $q^\mu \equiv (t, u, v)$ where t coincides with the laboratory time and u, v are the coordinates on the surface. Second, we have to use a Lagrangian rather than a Hamiltonian description. Consequently, the dynamics of graphene's conductivity electrons near a Dirac point is given by the customary generalization to a curved spacetime^{8,21} of the action for massless Dirac spinors in 2+1 dimensions¹⁵ (Supplementary Information I)

$$\mathcal{A} = i \int d^3q \sqrt{g} \bar{\psi}(q) \gamma^\mu \nabla_\mu \psi(q), \quad (2)$$

where $\hbar = v_F = k_B = 1$, with v_F the Fermi velocity.

We are building upon the continuum description of a Dirac quantum field, and upon modeling the effects of curvature through the coupling of the Dirac field to a curved *spatial* metric. The effectiveness of both assumptions was proved elsewhere^{22–24}. Here we assert that the electrons on graphene might directly experience a curved *spacetime* even though the curvature is all in the spatial part. That a spacetime can be curved acting solely on the spatial part is well known²⁵, the

breakthrough is to prove that this makes sense for graphene. In what follows we shall extract a measurable prediction based on such assumption and we shall show that the physical set-up to do that boils down to shape the graphene sample as a Beltrami pseudosphere²⁶.

When the metric (1) is conformally flat (Supplementary Information I), we can make use of the Weyl symmetry¹⁶ enjoyed by the action (2) to obtain exact results that otherwise are difficult or impossible to obtain. As proved in earlier work¹⁵ (see also Supplementary Information I), the metric (1) is conformally flat for all surfaces of constant Gaussian curvature $\mathcal{K}$, but we need (1) to *explicitly* take the form $g_{\mu\nu}^{\text{graphene}}(\mathcal{Q}) = \Phi^2(\mathcal{Q})g_{\mu\nu}^{\text{flat}}(\mathcal{Q})$, for certain coordinates $\mathcal{Q}^\mu$. Only then we can use all the power of Weyl symmetry. We know from general arguments that those coordinates must exist, but the key issue for graphene is whether such coordinates $\mathcal{Q}^\mu$ can be *practically realized in the laboratory*.

One of the main results of this work is that the Beltrami pseudosphere^{26,27}, $d\ell^2 = g_{\alpha\beta}q^\alpha q^\beta = du^2 + r^2 e^{2u/r} dv^2$, with $v \in [0, 2\pi]$, $u \in [-\infty, 0]$ (see Fig. 1), solves the problem. For this surface the coordinates $\mathcal{Q}^\mu$ are the coordinates q^μ , and the time coordinate coincides with the laboratory time. The isothermal coordinates are $\tilde{x} = v/r$ and $\tilde{y} = e^{-u/r}/r$ (Supplementary Information II), hence

$$ds_{\text{graphene}}^2 \equiv ds_{(B)}^2 = e^{2u/r} [e^{-2u/r} (dt^2 - du^2) - r^2 dv^2] . \quad (3)$$

As the line element in square brackets is flat (and Rindler^{8,10}, an exciting instance that we shall soon exploit), for a Beltrami pseudosphere we can fulfill the condition of a physically doable $g_{\mu\nu}^{\text{graphene}}(q) = \Phi^2(q)g_{\mu\nu}^{\text{flat}}(q)$ already in the frame q^μ . From now on, the spacetime we shall suppose

to be experienced by graphene's quasi-particles, is

$$g_{\mu\nu}^{(B)}(q) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -r^2 e^{2u/r} \end{pmatrix}, \quad (4)$$

where, $t \in [-\infty, +\infty]$, $u \in [-\infty, 0]$, $v \in [0, 2\pi]$. We call this a “Beltrami spacetime”.

Once we take this view, any measurement is an operation relating different spacetimes: the inner (2+1)-dimensional curved spacetime and the outer (3+1)-dimensional flat spacetime. See Fig. 2. The electrons, far from the graphene surface, loose the lattice-induced properties of quasi-particles, i.e. electric neutrality, masslessness, (2+1)-dimensionality and, the pseudo-relativistic Dirac nature. This latter fact, in particular, requires the final ground state reached after the measurement be of a non-relativistic nature. To accommodate this hybrid situation into the fully relativistic scenarios of QFT in curved spacetimes, when we take into account the different quantum vacua for the different observers involved^{8,9,11,28,29}, we approximate the ground state associated with the measurements with the Minkowskian $|0_M\rangle$. This way we have both, the mathematical description of the experimental evidences of the lattice-induced features (we keep a relativistic-like, (2+1)-dimensional structure), and the information on the true ground state (we use a flat vacuum defined everywhere). This is the best approximation: the Dirac field is living in a (2+1)-dimensional curved spacetime with coordinates $q^\mu \equiv (t, u, v)$, the measuring device has the same coordinates (i.e. it follows the profile of the surface in a specific manner described later), the quantum vacuum of reference entails information on the flatness of the ambient spacetime but retains the basic lattice-induced features.

3 Weyl symmetry and the fictitious Rindler spacetime

We shall focus on the one particle Green's function that contains all the information on the single particle properties of the system such as the LDOS, life time of the quasi-particles and thermodynamic properties (specific heat). For the reasons illustrated above, this is defined as

$$S^{(B)}(q_1, q_2) \equiv \langle 0_M | \psi^{(B)}(q_1) \bar{\psi}^{(B)}(q_2) | 0_M \rangle, \quad (5)$$

i.e. the positive frequency Wightman function, in the language of QFT on curved spacetimes^{8,10} (see also ²⁸). To obtain an exact result for this function, we shall use local Weyl symmetry^{15,16} as this case is a perfect match for its implementation: $g_{\mu\nu}^{(B)} = \varphi^2(u) g_{\mu\nu}^{(R)}$, $\psi^{(B)} = \varphi^{-1}(u) \psi^{(R)}$, with $\varphi(u) = e^{u/r}$ and the metric

$$g_{\mu\nu}^{(R)}(q) = \begin{pmatrix} e^{-2u/r} & 0 & 0 \\ 0 & -e^{-2u/r} & 0 \\ 0 & 0 & -r^2 \end{pmatrix} \quad (6)$$

describing a flat geometry. Local Weyl symmetry gives

$$\begin{aligned} \mathcal{A}_B &= i \int d^3q \sqrt{g^{(B)}} \bar{\psi}^{(B)}(q) \gamma^\mu \nabla_\mu \psi^{(B)}(q) \\ &= i \int d^3q \sqrt{g^{(R)}} \bar{\psi}^{(R)}(q) \gamma^\mu \nabla_\mu \psi^{(R)}(q) = A_R. \end{aligned} \quad (7)$$

Notice that $\mathcal{A}_B$ refers to a curved spacetime, whereas A_R to flat curvilinear coordinates. The Green's function of interest can then be written as

$$S^{(B)}(q_1, q_2) = \varphi^{-1}(q_1) \varphi^{-1}(q_2) S^{(R)}(q_1, q_2), \quad (8)$$

where

$$S^{(R)}(q_1, q_2) \equiv \langle 0_M | \psi^{(R)}(q_1) \bar{\psi}^{(R)}(q_2) | 0_M \rangle. \quad (9)$$

The superscript R here stands for “Rindler”. We shall now study the spacetime $g_{\mu\nu}^{(R)}$ to compute $S^{(R)}$, but one must bear in mind that this is a *fictitious* flat spacetime, the only *physical* spacetime being the (Weyl-equivalent) curved Beltrami spacetime $g_{\mu\nu}^{(B)}$. The Weyl transformation connects the Beltrami spacetime to a flat spacetime that, as well known, is the near-horizon spacetime for a black hole²¹. We claim that this setting is the closest possible to a black hole on graphene that can be obtained by solely acting on the spatial part of the metric. The physical Green’s function $S^{(B)}$ is recovered from $S^{(R)}$ through (8).

Let us introduce Minkowski coordinates $Q^\mu = (T, X, Y)$ for which $ds_{(R)}^2 = g_{\mu\nu}^{(R)} dq^\mu dq^\nu = \eta_{\mu\nu} dQ^\mu dQ^\nu$: $T = re^{-u/r} \sinh \frac{t}{r}$, $X = rv$, $Y = re^{-u/r} \cosh \frac{t}{r}$. Although no physical quantity explicitly depends from these coordinates, we have assumed that the quantum vacuum of reference is Minkowskian. We have that $Y^2 - T^2 = r^2 e^{-2u/r} \equiv \alpha^{-2}(u)$, which, for constant u , corresponds to worldlines of observers moving at constant “proper acceleration”^{8,10,21}

$$\alpha(u) \equiv \frac{e^{u/r}}{r} \in [0, r^{-1}]. \quad (10)$$

In a standard Rindler spacetime, the acceleration α can become infinite. In our case there is a maximal “acceleration”, $\alpha_{\max} \equiv \alpha(u = 0) = r^{-1} = \sqrt{-\mathcal{K}}$, of complete geometric origin. Furthermore, we are forever confined to one Rindler wedge ($\alpha \geq 0$) and on one side ($u \leq 0$) of the singular maximal circle (“Hilbert horizon”), the other side is inaccessible as “the world ends” at $R = r$ which corresponds to $u = 0$ (see $z(u)$ in the caption of Fig. 1). For a radius of curvature $r \sim 1\text{mm}$, the Rindler horizon $Y = T$ is effectively reached after a time $t_{\text{hor}} \sim r/v_F$ of few

nanoseconds ($v_F \sim 10^6 \text{m/s}$). Let us also introduce a perhaps more familiar notation

$$\eta = \frac{t}{r} = \alpha_{\max} t, \quad \xi = r e^{-u/r} = \alpha^{-1}(u), \quad (11)$$

with which $ds_{(R)}^2 = \xi^2 d\eta^2 - d\xi^2 - r^2 dv^2$.

The Rindler spacetime has been studied in great depth, many crucial results for massless fields are exact¹⁰, and is well known to give rise to a thermal spectrum for quantum fields¹⁸. Our task is to see whether those results can apply to graphene. The real coordinates q^μ are, through (11), those of an “accelerated” observer in the fictitious spacetime and constant “acceleration” means $u = \text{constant} \equiv \bar{u}$ (we also take $v = \text{constant} \equiv \bar{v}$). Thus, measuring at a fixed point $(\bar{u}, \bar{v})$ of the real Beltrami surface and letting time pass corresponds, in the equivalent fictitious Rindler spacetime, to measurements made by an observer on a worldline of constant proper “acceleration” $\alpha: t/r \equiv \alpha(\bar{u})\tau, u = \bar{u}, v = \bar{v}$.

To fit our set-up within the proper relativistic requirements we need then to refer to the Green’s function (9) *at the same point in space and at two different times*

$$S^{(R)}(t_1 - t_2, \mathbf{q}, \mathbf{q}) \equiv \langle 0_M | \psi^{(R)}(t_1; \mathbf{q}) \bar{\psi}^{(R)}(t_2; \mathbf{q}) | 0_M \rangle, \quad (12)$$

as this is what would be seen by an observer of the above described worldline. The dependence on $t_1 - t_2$ is a result of the stationarity of the worldline in point. Furthermore, we need to compute $S^{(R)}$ by setting $t \rightarrow t + i\varepsilon$ (where $t_1 - t_2 \equiv t$) a prescription that takes into account the nonzero size of the measuring device^{8,10}. For the Scanning Tunneling Microscopy (STM) experiment we have in mind, ε is the size, in “natural units”, of its needle or tip. For a tungsten needle $\varepsilon \sim$

$0.25\text{mm} \times v_F^{-1} \sim 10^{-10}\text{s}$, while for a typical tip $\varepsilon \sim 10\text{\AA} \times v_F^{-1} \sim 10^{-15}\text{s}$ (see, e.g.,³¹). Hence, the Unruh requests are satisfied by considering the Green's function $S^{(R)}(\tau, \mathbf{q}, \mathbf{q})$, where $\tau \equiv t/e^{\bar{u}/r}$, and measuring at each point for a time given by the largest among ε and the t_{hor} (that is $\sim 10^{-8}\text{s}$ for $r \sim 10\text{mm}$ and $\sim 10^{-12}\text{s}$ for $r \sim 1\mu\text{m}$). The Unruh thermal features are then readily seen by considering, as customary, the power spectrum^{8,10}

$$F^{(R)}(\omega, \mathbf{q}) \equiv \frac{1}{2} \text{Tr} \left[\gamma^0 \int_{-\infty}^{+\infty} d\tau e^{-i\omega\tau} S^{(R)}(\tau, \mathbf{q}, \mathbf{q}) \right], \quad (13)$$

that, for graphene, besides inessential constants, coincides with the definition of the electronic LDOS^{24,32,33}, $\rho^{(R)}(\omega, \mathbf{q}) \equiv \frac{2}{\pi} F^{(R)}(\omega, \mathbf{q})$. As we are in a massless model, the quantity in (13) can be computed exactly to give¹⁰ $F^{(R)}(\omega, \mathbf{q}) = \frac{1}{2} \omega / (e^{\omega/\mathcal{T}} - 1)$, where $\mathcal{T}$ is a Unruh temperature¹⁸

$$\mathcal{T} \equiv \frac{\alpha(\bar{u})}{2\pi} = \frac{e^{\bar{u}/r}}{2\pi r} \equiv \mathcal{T}_0 e^{\bar{u}/r}, \quad (14)$$

that includes the constant $\mathcal{T}_0 = 1/(2\pi r)$ and the Tolman factor²¹ $e^{\bar{u}/r}$, required by the local measurements. We see here the appearance of a thermal distribution, the typical effect predicted by QFT in this context. That is a Bose-Einstein distribution for a Dirac field, an instance theoretically predicted to be a general feature (“swapping of statistics”) of both the Hawking and Unruh phenomena in odd dimensional spacetimes^{10,19,20}. Nonetheless, as said in various occasions, the only physical set-up we have is the Beltrami spacetime. To discuss the physics we have now to move to that case.

4 Experimental prediction of the physical LDOS

The task is easy due to Weyl symmetry. In fact, we have tuned from the very beginning the whole procedure towards the optimal use of this symmetry. The Weyl factor in (8) being time-independent, goes through the Fourier transform in (13), hence the LDOS we predict for a graphene sheet shaped as a Beltrami pseudosphere is $\rho^{(B)}(\omega, \mathbf{q}) = \varphi^{-2}(\mathbf{q})\rho^{(R)}(\omega, \mathbf{q})$, or, more explicitly

$$\rho^{(B)}(E, \bar{u}, r) = \frac{4}{\pi} \frac{1}{(\hbar v_F)^2} \frac{E e^{-2\bar{u}/r}}{\exp[E/(k_B \mathcal{T}_0 e^{\bar{u}/r})] - 1}, \quad (15)$$

where we included the $g = 4$ degeneracy, and the proper dimensional units are re-introduced, for instance $\omega \equiv \omega/v_F$, $E \equiv \hbar\omega$ and $\mathcal{T}_0 \equiv \hbar v_F/(k_B 2\pi r)$. This equation is our prediction of the exact functional dependence of the LDOS from the energy E , the meridian coordinate u and the curvature r (even if we explicitly wrote-in $\mathcal{T}_0$ this is essentially $\sim r^{-1}$). With the notation $\bar{u}$ we want to stress that: i) the measuring device has to closely follow the profile of the surface, so that its coordinates can be taken to be q^μ ; and ii) the device has to stop at each given point on the surface $(\bar{u}, \bar{v})$ for a time much bigger than $\max(\varepsilon, t_{\text{hor}})$, not a stringent condition, as we saw earlier.

We shall soon describe how to detect this Hawking-Unruh effect through an STM experiment, but let us first make some comments. We do not have a black-hole here, but we do not need one as our Hawking effect is inferred from an equivalent (through Weyl symmetry) Unruh phenomenon. Similar in spirit are the well known results of ³⁴, where the Hawking effect is obtained from a higher dimensional Unruh effect. Our results differ from those because we ruled-out the embedding into un-physical higher dimensional spacetimes and because the Beltrami spacetime is not a proper anti de Sitter (AdS) (as a consequence, our results are closer to the dS case of ³⁴ rather

than its AdS).

Let us focus now on the experimental detection. To realize with an STM device in the lab the situation depicted in Fig. 2 we need to follow the prescriptions i) and ii) above and we have to make sure that the tunneling current measured is the one relative to the sample electrons tunneling to the tip, and not viceversa. This is easily achieved by fixing the right polarity of the bias voltage. The plot we need is a current-to-voltage (dI/dV) spectrum. This is obtained by first constructing a one-layer graphene sample shaped as a given Beltrami pseudosphere with fixed r and by measuring at a particular fixed value of u . We then vary the bias voltage to obtain the $\rho^{(B)}$ vs E behavior. As the formula (15) does not depend on the angle v , we do not need to stay at a fixed point (u, v) on the surface but we can loop around the circle at fixed u (see Fig. 1) while varying the bias voltage V . We can then repeat the same measurements at different values of u to obtain a series of dI/dV spectra (a “line-cut”). The expected line-cut is illustrated in Fig. 3 for $r = 10\mu\text{m}$.

To avoid defects proliferation we need to take large curvature on the natural scale furnished by the lattice spacing $\ell \sim \text{\AA}$: $r \gg \ell$. We take $r \sim 10^{-7}\text{m}$ as the highest curvature. This corresponds to $\mathcal{T}_0 \sim 13\text{K}$ as the highest detectable temperature (this value can only be made smaller by including the effects of the local measurement, $e^{\bar{u}/r}$, as $u < 0$). The zero-curvature limit (large r) of $\rho^{(B)}$ does not match the flat LDOS, $\rho^{(\text{flat})}(E) = \frac{2}{\pi} \frac{1}{(\hbar v_F)^2} |E|$. This is as it must be since (15) is not the result of a perturbative computation with $\rho^{(\text{flat})}(E)$ as the leading term, thus even a very small curvature fully turns on the effect. In Fig. 4 we show the behavior of $\rho^{(B)}$ for a wide range of Beltrami pseudospheres with curvatures spanning a range of 6 orders of magnitude.

As can be seen there, fixing the value of u in units of r , the effect is the same for all curvatures.

The limit for zero energy gives a nonzero result, $\rho^{(B)}(0, \bar{u}, r) = \frac{2}{\pi^2} \frac{1}{\hbar v_F} \frac{e^{-\bar{u}/r}}{r}$, as expected for a finite temperature LDOS. A direct measurement of this value of the LDOS would give the constant Hawking temperature $\mathcal{T}_0 = \frac{\pi}{4} \frac{\hbar v_F}{k_B} e^{\bar{u}/r} \rho^{(B)}(0, \bar{u}, r)$, nonetheless, this measurement seems difficult to perform due to the extremely high sensitivity necessary for the appreciation of this effect, see inset of Fig. 4. It would be difficult to realize whether it is the effect of a finite temperature or, e.g., of impurities.

The best way to measure the Hawking temperature is the one based on the line-cut in Fig. 3 and described above. For each power spectrum (fixed u) we shall obtain the corresponding u -dependent temperature, $\mathcal{T} = E/[k_B \ln(\frac{4Ee^{-2\bar{u}/r}}{\pi(\hbar v_F)^2 \rho^{(B)}} + 1)]$. Different $\mathcal{T}$ s for different u s will then give the same constant Hawking temperature through $\mathcal{T}_0 = e^{-u/r} \mathcal{T}$.

1. Wallace, P. R. The band theory of graphite. *Phys. Rev.* **71**, 622-634 (1947).
2. Semenoff, G. W. Condensed-matter simulation of a three-dimensional anomaly. *Phys. Rev. Lett.* **53**, 2449-2452 (1984).
3. Novoselov, K.S. *et al.* Electric field effect in atomically thin carbon films. *Science* **306** 666-669 (2004).
4. Novoselov, K.S. *et al.* Two-dimensional gas of massless Dirac fermions in graphene. *Nature* **438** 197-200 (2005).

5. Geim, A. K. Graphene: status and prospects. *Science* **324** 1530-1534 (2009).
6. Volovik, G. E. *The universe in a helium droplet* (Clarendon Press, Oxford, 2003).
7. Novello, M., Visser, M. and Volovik, G. E. *Artificial Black Holes* (World Scientific, Singapore, 2002).
8. Birrell, N. D. and Davies, P. C. W. *Quantum fields in curved space* (Cambridge Univ. Press, Cambridge, 1982).
9. Wald, R. M. *Quantum field theory in curved spacetimes and black hole thermodynamics* (The Univ. Chicago Press, Chicago, 1994).
10. Takagi, S. Vacuum noise and stress induced by uniform acceleration. *Prog. Theor. Phys. Suppl.* **88** 1-142 (1986).
11. Fulling, S. A. and Ruijsenaars, S. N. M. Temperature, periodicity and horizons. *Phys. Rept.* **152** 135-176 (1987).
12. Deser, S., Jackiw, R. and 't Hooft, G. Three-dimensional gravity: dynamics of flat space. *Ann. Phys.* **152** 220-235 (1984).
13. Carlip, S. *Quantum gravity in 2+1 dimensions* (Cambridge Univ. Press, Cambridge, 1998).
14. Castro Neto, A. H. *et al.* The electronic properties of graphene. *Rev. Mod. Phys.* **81** 109-162 (2009).
15. Iorio, A. Weyl-gauge symmetry of graphene. *Ann. Phys.* **326** 1334-1353 (2011).

16. Iorio, A., O’Raifeartaigh, L., Sachs, I. and Wiesendanger, C. Weyl gauging and conformal invariance. *Nucl. Phys. B* **495** 433-450 (1997).
17. Hawking, S. Black hole explosion. *Nature* **248** 30-31 (1974).
18. Unruh, W. G. Notes on black-hole evaporation. *Phys. Rev. D* **14** 870-892 (1976).
19. Hyun, S., Lee, G. H. Yee, J. H. Hawking radiation from a $(2 + 1)$ -dimensional black hole. *Phys. Lett. B* **322** 182-187 (1994).
20. Hyun, S., Lee, G. H. Yee, J. H. Hawking radiation of Dirac fields in the $(2 + 1)$ -dimensional black hole spacetime. **51** 1787-1792 (1995).
21. Wald, R. M. *General relativity* (The Univ. Chicago Press, Chicago, 1984).
22. Gonzalez, J., Guinea, F. and Vozmediano, M. A. H. Continuum approximation to fullerene molecules. *Phys. Rev. Lett.* **69** 172-175 (1992).
23. Gonzalez, J., Guinea, F. and Vozmediano, M. A. H. The electronic spectrum of fullerenes from the Dirac equation. *Nucl. Phys. B* **406** 771-794 (1993).
24. de Juan, F., Cortijo, A. and Vozmediano, M. A. H. Charge inhomogeneities due to smooth ripples in graphene sheets. *Phys. Rev. B* **76** 165409-1-165409-9 (2007).
25. Feynman, R. P., Morinigo, F. B., and Wagner, W. G., *Feynman’s lectures on gravitation* (Addison-Wesley, Reading MA, 1995).
26. Eisenhart, L. P. *A treatise on the differential geometry of curves and surfaces* (Princeton Univ. Press, Princeton, 1940).

27. Spivak, M. *A comprehensive introduction to differential geometry, part III* (Publish or Perish Inc., Houston, 1999).
28. Iorio, A., Lambiase, G. and Vitiello, G. Entangled quantum fields near the vent horizon and entropy. *Ann. Phys.* **309** 151-165 (2004).
29. Israel, W. Thermo-field dynamics of black holes. *Phys. Lett. A* **57** 107-110 (1976).
30. Hilbert, D. and Cohn-Vossen, S. *Geometry and the imagination* (Chelsea Pub., New York, 1952).
31. Tung, F., Yoshimura, M. and Ueda, K. Direct fabrication of Carbon nanotubes STM tips, *J. Nanomat.* **2009** 612549 (2009).
32. Rickayzen, G. *Green's functions and condensed matter* (Academic Press, London, 1980).
33. Altland, A. and Simons, B. *Condensed matter field theory* (Cambridge Univ. Press, Cambridge, 2006).
34. Deser, S. and Orit Levin, Accelerated detectors and temperature in (anti-) de Sitter spaces, *Class. Quant. Grav.* **14** L163-L168 (1997).
35. Kleinert, H. *Gauge fields in condensed matter. Vol II* (World Scientific, Singapore, 1989).
36. Katanaev, M. O. and Volovich, I. V. Theory of defects in solids and three-dimensional gravity. *Ann. Phys.* **216** 1-28 (1992).
37. Vozmediano, M. A. H., de Juan, F. and Cortijo, A. Gauge fields and curvature in graphene. *J. Phys. Conf. Series* **129** 012001 (2008).

SUPPLEMENTARY INFORMATION

The Hawking-Unruh phenomenon on graphene

A. Iorio, G. Lambiase

I. Graphene Dirac spinors and geometric conventions. Due to the special geometry of graphene's Honeycomb lattice (two interpenetrating triangular lattices), the Bloch wave function has two components representing the probability amplitude at each sublattice [S1]. The spinorial nature of the excitations is linked to this sublattice degree of freedom named pseudospin in the graphene literature. The other important ingredient of the Dirac model, also a consequence of the special topology of the lattice, is the fact that the tight binding dispersion relation at half filling has six Fermi points instead of a Fermi line [S1]. As only two of them are inequivalent this introduces an extra degeneracy of the vacuum. This new degree of freedom is called valley in semiconductor language and plays an important role in the physics of graphene [S1]. Linearizing around the two inequivalent Fermi points (Dirac points), $\vec{k}_\pm^D$, $\vec{k}_\pm \simeq \vec{k}_\pm^D + \vec{p}$, the tight-binding Hamiltonian can be written as ($\hbar = 1$) [S1]

$$\begin{aligned} H &= v_F \sum_{\vec{p}} \left(\psi_+^\dagger \vec{\sigma} \cdot \vec{p} \psi_+ + \psi_-^\dagger \vec{\sigma}^* \cdot \vec{p} \psi_- \right) \\ &= -iv_F \int d^2x \left(\psi_+^\dagger \vec{\sigma} \cdot \vec{\partial} \psi_+ + \psi_-^\dagger \vec{\sigma}^* \cdot \vec{\partial} \psi_- \right), \end{aligned} \quad (\text{S1})$$

with $\vec{\sigma} \equiv (\sigma_1, \sigma_2)$, $\vec{\sigma}^* \equiv (-\sigma_1, \sigma_2)$, σ_i being the Pauli matrices, the Fermi velocity $v_F \equiv 3t\ell/2$ (that will be set to 1) $t \simeq 2.7$ eV is the hopping parameter, $\ell \simeq 2.5\text{\AA}$ is the lattice spacing. Where $\psi_+^T \equiv (a_+ \ b_+)$, $\psi_-^T \equiv (a_- \ b_-)$ are two-component Dirac spinors, as appropriate for this

2+1-dimensional system, and a and b are anti-commuting annihilation operators for an electron in the two sub-lattices. In here, we do not consider short range scattering centers or any other effect mixing the two Fermi points, thus we discuss the physics around a *single* Fermi point, e.g. $\psi \equiv \psi_+$. The corresponding action is $A = i \int d^3x \bar{\psi} \gamma^a \partial_a \psi$, where $\gamma^0 = \sigma_3$, $\gamma^1 = i\sigma_2$, $\gamma^2 = -i\sigma_1$ which obey $[\gamma^a, \gamma^b]_+ = 2\eta^{ab}$, with $a, b = 0, 1, 2$ the Lorentz/flat indices (see below). The curved action is

$$\mathcal{A} = i \int d^3x \sqrt{g} \bar{\psi} \gamma^a E_a^\mu \nabla_\mu \psi, \quad (\text{S2})$$

where we implicitly introduced the Vielbein e_μ^a and its inverse E_a^μ , $\eta_{ab} e_\mu^a e_\nu^b = g_{\mu\nu}$, $e_\mu^a E_a^\nu = \delta_\mu^\nu$, $e_\mu^a E_b^\mu = \delta_b^a$, the indices $\mu, \nu = 0, 1, 2$ respond to diffeomorphisms (Einstein indices), while $a, b = 0, 1, 2$ respond to flat space transformations (Lorentz indices), $\eta_{ab} = \text{diag}(+1, -1, -1)$, $\sqrt{g} = \sqrt{\det g_{\mu\nu}}$ and the diffeomorphic covariant derivative is $\nabla_\mu = (\partial_\mu + \frac{1}{2} \omega_\mu^{bc} J_{bc})$, with $J^{ab} = \frac{1}{4} [\gamma^a, \gamma^b]$, and $\omega_\mu^a{}_b = e_\lambda^a (\delta_\nu^\lambda \partial_\mu + \Gamma_{\mu\nu}^\lambda) E_b^\nu$ is the spin connection obtained by requiring the full covariant derivative of the Vielbein to be zero (metricity condition) $\nabla_\mu e_\nu^a = \partial_\mu e_\nu^a - \Gamma_{\mu\nu}^\lambda e_\lambda^a + \omega_\mu^a{}_b e_\nu^b = 0$, where $\Gamma_{\mu\nu}^\lambda$ is the Christoffel connection. Our convention for the Riemann curvature tensor is $R^\rho{}_{\lambda\mu\nu} = \partial_{[\nu} \Gamma^\rho{}_{\mu]\lambda} + \Gamma^\rho{}_{[\nu\sigma} \Gamma^\sigma{}_{\mu]\lambda}$, that, in terms of $\omega_\mu^a{}_b$ is $(R_{\mu\nu})^a{}_b = (\partial_{[\nu} \omega_{\mu]}^a{}_b + \omega_{[\nu}^a{}_c \omega_{\mu]}^c{}_b)$. Torsion $T^\lambda{}_{\mu\nu} = \Gamma^\lambda{}_{[\mu\nu]}$ is taken to be zero.

For the metric (1) the Ricci tensor is $R_\mu{}^\nu = \text{diag}(0, \mathcal{K}, \mathcal{K})$ which gives as the only nonzero components of the Cotton tensor, $C^{\mu\nu} = \epsilon^{\mu\sigma\rho} \nabla_\sigma R_\rho{}^\nu + \mu \leftrightarrow \nu$, the result $C^{0x} = -\partial_y \mathcal{K} = C^{x0}$ and $C^{0y} = \partial_x \mathcal{K} = C^{y0}$. Since conformal flatness in (2+1)-dimensions amounts to $C^{\mu\nu} = 0$, this shows that all surfaces of constant $\mathcal{K}$ give rise in (1) to conformally flat 2+1 dimensional spacetimes. Note that the result holds for $(+, -, -)$ and for $(+, +, +)$.

II. Beltrami spacetime and physical coordinates Q^μ . We found that the coordinates Q^μ for the sphere, the first example one would consider, are not easy to envisage, while surfaces with constant negative curvature, $\mathcal{K} < 0$, are a better candidate. For the latter case the metric of graphene can be written in the spatial isothermal coordinates [S2], i.e., in terms of the line element, as $ds_{\text{graphene}}^2 = dt^2 - \frac{r^2}{\tilde{y}^2}(d\tilde{x}^2 + d\tilde{y}^2)$ where $\tilde{x}, \tilde{y}$ are the *abstract* coordinates of the Lobachevsky geometry in the upper half-plane ($\tilde{y} > 0$) model and $r = \sqrt{-\mathcal{K}^{-1}}$ [S3]. The actual type of surface is obtained by specifying $\tilde{x}$ and $\tilde{y}$ in terms of coordinates measurable using the Euclidean distance (embedding). This is crucial for the metric to describe the spacetime experienced by the electrons on real graphene (an instance not considered in [S4,S5]).

The key fact common to all these surfaces (whether or not they are surfaces of revolution) is a theorem by Hilbert that proves *the impossibility to have a complete surface of constant negative Gaussian curvature embedded in $\mathbb{R}^3$* [S6]. Complete means, essentially, that there are no singularities (if one changes the dimensions, e.g., $\mathbb{R}^3 \rightarrow \mathbb{R}^4$, or the signature, $(+++)$ $\rightarrow$ $(-++)$, the task becomes indeed possible), hence we must expect singularities of some sort in our surfaces. By writing the line element as $ds_{\text{graphene}}^2 = \frac{r^2}{\tilde{y}^2} \left[\frac{\tilde{y}^2}{r^2} dt^2 - d\tilde{x}^2 - d\tilde{y}^2 \right]$ it is easy to check that the line element in square brackets is flat. This apparently solves our problem as the Q^μ that should make sense appear to be $(t, \tilde{x}, \tilde{y})$. But it is not so until we identify the specific surface and reduce to the measurable coordinates. For instance, we found that the elliptic pseudosphere [S6], $d\ell^2 = du^2 + r^2 \sinh^2 u/r dv^2$ (here u and v are the meridian and parallel coordinate on the surface, respectively) has as natural coordinate system a different one, namely $(T = e^{t/r} \cosh u/r, X = e^{t/r} \sinh u/r \cos v, Y = e^{t/r} \sinh u/r \sin v)$. Clearly, it is very difficult to

imagine how to practically realize a physical situation in the laboratory so that those coordinates become meaningful. It should be now evident that the identification of the right surface is quite an art.

S1. Castro Neto, A. H. *et al.* The electronic properties of graphene. *Rev. Mod. Phys.* **81** 109-162 (2009).

S2. Chern, S. An elementary proof of the existence of isothermal parameters on a surface. *Proc. Amer. Math. Soc.* **6** 771-782 (1955).

S3. Penrose, R. *The road to reality* (Knopf, New York, 2005).

S4. Gorbar, E. V. & Gusynin, V. P. Gap generation for Dirac fermions on Lobachevsky plane in magnetic field. *Ann. Phys.* **323** 2132-2146 (2008).

S5. Boada, O., Celi, A., Latorre, J. I. & Lewenstein, M. Dirac equation for cold atoms in artificial curved spacetime. *New J. Phys.* **13** 035002 (23pgs) 2011.

S6. Hitchin, N. *Geometry of surfaces* Lecture Notes Oxford (Oxford) 2004.

Acknowledgements We are indebted to Maria A.H. Vozmediano for providing guidance in crucial condensed matter issues and for many enlightening discussions. We also thank I. Brihuega, F. Guinea and N. Hitchin for correspondence. A. I. thanks S. Bonanos for providing free software for geometric tensors' computation and G. L. acknowledges the financial support of MIUR through PRIN (Prot. 2008NR3EBK



002).

Competing Interests The authors declare that they have no competing financial interests.

Correspondence Correspondence and requests for materials should be addressed to

A. Iorio (email: alfred.iorio@gmail.com).

Author contributions All authors contributed extensively to the work presented in this paper.







Figure 1 Beltrami pseudosphere. The $\mathbb{R}^3$ coordinates of the Beltrami pseudosphere, in the canonical form, $(R(u) \cos v, R(u) \sin v, z(u))$, are $x(u, v) = c e^{u/r} \cos v$, $y(u, v) = c e^{u/r} \sin v$, $z(u) = r(\sqrt{1 - \frac{c^2}{r^2} e^{2u/r}} - \operatorname{arctanh} \sqrt{1 - \frac{c^2}{r^2} e^{2u/r}})$, $c > 0$ and $r = \sqrt{-\mathcal{K}^{-1}} > 0$ with $\mathcal{K}$ the Gaussian curvature. We choose $c = r$, thus $R(u) = r e^{u/r} \in [0, r]$ as $u \in [-\infty, 0]$. The curves at constant angle v are the meridian curves and coincide with the null geodesics of the spacetime ($d\ell^2 = dt^2$).

Figure 2 Choice of vacuum. Graphene quasi-particles enter the measuring device as a (2+1)-dimensional massless Dirac “curved” field $\psi^{(B)}$ (a). Eventually, they loose the lattice-induced features, and behave as charged, massive, non-relativistic electrons in a (3+1)-dimensional spacetime (c). To include both, the routinely detected lattice-induced features, and the flat status of the ground state eventually reached, the Green’s functions need to be expectations of products of $\psi^{(B)}$ fields on a flat (2+1)-dimensional vacuum $|0_M\rangle$ (b). The Minkowski vacuum, being defined everywhere, mimics the true vacuum $|0_E\rangle$ better than the Rindler vacuum $|0_R\rangle$ that only lives in one sector of the spacetime.

Figure 3 Line cut. Series of dI/dV spectra (“line-cut”) obtainable with an STM. In this example the curvature is $r = 10\mu\text{m}$. The measurements need to be performed at a fixed value of u , but since $\rho^{(B)}$ does not depend on the angle v , we can loop around the circle at fixed u while varying the bias voltage V . We can then repeat the same measurements at different values of u to obtain the shown line-cut. The polarity of the bias V is such that the sample electrons tunnel to the STM tip, and not viceversa (see Fig. 2). The

energy range is $E \in [-1, 1]\text{eV}$, where the linear behavior of the flat LDOS (shown in blue) is more pronounced. The range of the meridian coordinate is $u \in]0, -3r]$. The $u = 0$ value (red dashed curve) corresponds to the singular boundary of the pseudosphere, hence we shall not trust our model there. The $u = -3r$ value (green curve) is chosen because beyond that value the pseudosphere is too sharp (line shaped, see Fig. 1). To each spectrum corresponds a u -dependent temperature $\mathcal{T}$ that ranges from $\sim 0.05\text{mK}$ till $\sim 0.1\text{K}$. The constant temperature (same for the whole line-cut) is then obtained through $\mathcal{T}_0 = e^{-u/r}\mathcal{T} \sim 0.13\text{K}$.

Figure 4 Nonperturbative effect. dI/dV spectra obtainable with an STM for different Beltrami pseudospheres (different rs). The main plot shows that the effect is present for a wide range of curvatures, from very large to very small $r \in [10^{-7}, 10^{-2}]\text{m}$. Here the value of the meridian coordinate is chosen to be $u = -r$ for all curvatures, giving rise to the same curve for them all. The inset shows the near zero behavior of $\rho^{(\text{B})}$ for the highest curvature considered here, $r = 10^{-7}\text{m}$, that is the most favorable for the experimental detection of a finite temperature at $E = 0$. The intercept of dashed lines with the y -axis gives the $\rho^{(\text{B})}(E = 0)$ values. Despite of that the required instrument sensitivity for such detection is too high (the axes of the inset have the same units as for the main plot), and it would not be clear whether it is the effect of a finite temperature or, e.g., of impurities.